



Phytotherapeutic, Nutraceutical, Medicinal, and Forage Properties of Fenugreek (*Trigonella foenum-graecum* L.): A Comprehensive Review

Sharda Choudhary ^a, Ravindra Singh ^{a*}, Ravi, Y ^a,
CB Gena ^b, Dheeraj Singh ^c, CB Harisha ^d, RS Meena ^a,
NK Meena ^a, VBS Chauhan ^e, Kavan Kumar V ^f,
Versha Kumari ^g and Arvind Kumar Verma ^a

^a ICAR- National Research Centre on Seed Spices, Ajmer- 305 206, Rajasthan, India.

^b Maharaja Ganga Singh University, Bikaner, Rajasthan Trustee,
Krishan-Gopal Ayurved Sansthan, Kaleda, Ajmer- 305 201, Rajasthan, India.

^c ICAR-Central Arid Zone Research Institute, Jodhpur – 342 003, Rajasthan, India.

^d ICAR-National Institute of Abiotic Stress Management, Baramati-413115, Maharashtra, India.

^e Regional Centre, ICAR- Central Tuber Crops Research Institute, Bhubaneswar -751 019, Odisha, India.

^f Department of Renewable Energy Engineering, CTAE, MPUAT, Udaipur- 313 001, India.

^g Sri Karan Narendra Agriculture University, Jobner- 303 329, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author RS conceptualized the topic. Authors SC and RY collected the literature and wrote the manuscript. Authors RS, SC, RY, CBG and DS checked and edited the manuscript. All the authors read and approved the manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i101494>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123839>

Review Article

Received: 25/07/2024
Accepted: 27/09/2024
Published: 03/10/2024

*Corresponding author: E-mail: mahla_rs@yahoo.com;

Cite as: Choudhary, Sharda, Ravindra Singh, Ravi, Y, CB Gena, Dheeraj Singh, CB Harisha, RS Meena, NK Meena, VBS Chauhan, Kavan Kumar V, Versha Kumari, and Arvind Kumar Verma. 2024. "Phytotherapeutic, Nutraceutical, Medicinal, and Forage Properties of Fenugreek (*Trigonella Foenum-Graecum* L.): A Comprehensive Review". *Journal of Advances in Biology & Biotechnology* 27 (10):719-33. <https://doi.org/10.9734/jabb/2024/v27i101494>.

ABSTRACT

Trigonella foenum-graceum L., commonly known as Fenugreek, is a yearly plant from the Fabaceae family. It boasts a wide array of uses and economic importance in both domestic and global markets. In India, it is grown primarily as a winter crop across several states, including Rajasthan, Madhya Pradesh, Gujarat, Andhra Pradesh, Chhattisgarh, Haryana, Punjab, Telangana, and Uttaranchal. Historically, Fenugreek has played a significant role in Ayurveda, an ancient Indian medicinal tradition, for treating various ailments in both humans and animals. Its leaves are often eaten as a vegetable and are used to prepare dishes like "parathas" and various snacks such as Biscuits, Laddoo, Tadka, and Sprouts. In regions like Rajasthan and Punjab, Fenugreek is primarily grown for its use as green or dry fodder for livestock. As a plant in the Fabaceae family, Fenugreek's root nodules are responsible for converting atmospheric nitrogen into nitrate, which enhances nitrogen fixation and improves soil health. Fenugreek seeds and leaves are known for their diverse health benefits, including anti-diabetic, anti-cancer, anti-inflammatory, antioxidant, anti-ulcer, anti-fertility, and immune-modulating properties. Presence of 4-hydroxyisoleucine and diosgenin in fenugreek seed has several beneficial effects and gaining popularity in ayurveda. It contains high fiber, protein and several bioactive compounds which make fenugreek a natural and health-promoting herb.

Keywords: *Economic importance; ethnobotany fenugreek; phytotherapeutics; phytonutraceutical; Trigonella foenum-graceum.*

1. INTRODUCTION

According to a World Health Organization (WHO) report, out of more than 400 families of medicinal plants, 315 occur in India. Thus, the availability of traditional knowledge and resources makes India a potential leader in the future pharmaceutical industry [1]. There is an urgent need that the knowledge of Charaka (1st century BCE), Sushruta (6th Century BCE), Vagbhata (6th century AD) and other Ayurvedic seers be tapped and delivered to the populace using the productivization methodology currently available.

In India, fenugreek (*Trigonella foenum-graecum* Linn.), popularly known as "methi" is a legume seed spice and medicinal herb belonging to the family Fabaceae. The fenugreek plant is an erect annual herb. *Trigonella* signifies 'little triangle' because of its yellowish-white triangular blossoms [2]. Some common regional names of this plant are hayseed (English), Hulba (Arabic), Moshoseitaro (Greek), Shoot (Hebrew) and mukhme shampeet (Persian), methika (Sanskrit), methiya (Kannad), menticura (Telugu) Uluva (Malayalam). Fenugreek is one of the most established winter crops under cultivation in Asia 4000 BC. It has been postulated that its centre of origin could be the Indian sub-continent as it is broadly cultivated in India and other neighbouring Asian nations [3]. In India, fenugreek is cultivated for human consumption as well as forage, fodder, seed and green manure [4,5]. Dry seeds of fenugreek have a strong aroma and slight

bitterness in taste. These are utilized as flavours, seasoning, preservatives, spice and flavouring agents for pickles, sprouts, curry, and vegetables to improve the flavour, aroma and nutritive value. These seeds are rich in fiber, gum and other phytonutrients. Extracts from fenugreek seed include diosgenin, saponin and have phytotherapeutic potential [6]. Fenugreek can potentially suppress microbial infections [7,8] in companion crops, humans, and animals. It has additionally been assessed for green manure for its nitrogen-fixing potential [9,10]. The high fiber, gum and protein present in seeds are used as a food stabilizer, additive and emulsifier. Extracts of fenugreek seed containing galactomannan are utilized as nutrients, industrial thickeners and therapeutic purposes [11,12,5]. Fenugreek is an essential dietary part of Indian flavours and is utilized to improve health because of the expectancy of medical advantages. Fenugreek is notable for using in nourishment to forestall ailments such as intestinal gas, cramps, swelling, acid reflux, diabetes, malignant growth, hypertension, joint inflammation, liver problems and numerous different illnesses and disorders. Because of the significant levels of cell reinforcement mixes, antioxidants, bioactive compounds, phenols, flavonoids, and anthocyanin, different tests and clinical preliminaries have demonstrated that fenugreek can fight pathologic conditions, particularly for the treatment and avoidance of perilous sicknesses, for example, diabetes, few malignant growths, contaminations and gastrointestinal disorders.

To utilize the medicinal properties of fenugreek at its full potential, demand is increasing over time to develop the products with new clinical proofs. Fenugreek is treated as a decent contender for traditional and ayurvedic drugs as it has phenolic compounds, bioactive amino acids, glycosides and cancer prevention agent activities. Despite the unique logical and clinical profile of fenugreek, consumer understanding is still in its adolescence and growers/promoters of this spice need to concentrate on this significant economic and commercial obstacle. However, in recent years, a good number of research and publications have been published on the yield and economics, fertility and biofertilizer effect on quality and productivity [13], nutritional value and uses [14], fodder bank [15], protein structure [16], molecular characterization [17] and organic production [18] of fenugreek to understand its more roles in medical uses.

Agrobiodiversity occupies a unique place within the overall realm of biodiversity. It refers to diversity among crops, cultivation systems, agroecosystems, horticulture, animals, birds, insects and microorganisms. Plant genetic diversity within cultivated crops is a key component of agricultural production systems. To ensure sustainable agriculture in diverse agroecosystems while implementing good agricultural practices and new livestock systems to intensify agriculture and alleviate increasing

pressures on natural ecosystems. Conserving naturally available resources and plant diversity should be a priority. Traditional crops such as fenugreek have the potential to be conserved if they are tied to the economic development of farmers [19].

2. BOTANY AND MORPHOLOGY

Fenugreek is an annual herb with $2n = 16$ chromosome numbers. The plant is an erect, hairy, cylindrical, branched medium height (30-80 cm) and has root nodules on branched tap root. The pod of this plant contain around 16 to 20 tiny, yellowish-dark coloured, sweet-smelling, and impactful seeds (Fig. 1). Leaves comprise three-lobed contrarily applaud pamphlets with short petioles, serrated edges and oval stipules [20]. Flowers; 1-2 axillary, sessile, bracteate, zygomorphic, corolla much exerted, white or creamy-white. Flowers have five petals, known as the banner, wing, and keel. It is a self-pollinating crop. Fenugreek seeds are yellow to greenish-brown, rhomboidal, flat, 5-8mm long with a deep groove across one corner. It is cultivated as a cold-season seed spice and fodder crop, mainly in north-western India. It is grown during winter (October to April) and flowers during January-February and March-April. Its leaves are used mainly as vegetables, fodder and seeds for pot-herb medicinal properties.




Fig. 1. Fenugreek (*Trigonella foenum-graecum* Linn.) (A) Field view of fenugreek crop at ICAR-NRCCS, Ajmer, (B) Bunch of leaves, (C) Flowering stage, (D) Pod in fenugreek, (E) Root nodules of fenugreek, (F) Mature seed, (G-K) Products of fenugreek: (G) Salty Biscuits, (H) Sweet Biscuits, (I) Laddoo, (J) Daal tadka and (K) Sprouts

3. NUTRITIONAL CONSTITUTES OF HERB

Fenugreek's high nutritional value makes it suitable for various applications, including vegetables, food, beverages, nutraceuticals, pharmaceuticals, flavour and fragrance, and several industrial uses. Trigonelline, nicotinic acid, and other alkaloids are found in the stem etc [21]. Rao [22] reported that fenugreek leaves contain seven saponins known as lectins, which are chemically glycosides of diosgenin. Leaves contain approximately 86.1% water, protein (4.4%), fat (0.9%), minerals (1.5%), fiber (1.1%), and sugar (6%). Leaves are a rich source of minerals such as calcium, phosphorus, magnesium, zinc, iron, and major amino acids such as riboflavin, thiamine and niacin. It was found that crunchy fenugreek herb contained 220.97 mg of ascorbic acid per 100 g of leaves and β -carotene 19 mg/100 g. Young leaves have excellent nutrient retention, so eating them raw or cooked can help you lose weight [23].

4. QUALITY PARAMETERS OF FENUGREEK SEEDS

Fenugreek seeds are the most valuable part of the plant. The seeds are generally golden yellow and have a maple flavour but a bitter taste. Seeds are fibrous, sticky and rubbery [24]. Fenugreek contains many chemical compounds, including steroidal saponins. A diosgenin

component was found in the oily germ of fenugreek [25]. Table 1 depicts the nutritional and pharmaceutical components of fenugreek.

5. CONSTITUENTS OF SEED AND OIL

Seeds are a rich source of gum, oil, amino acids, tryptophan and lysine, dietary fiber, alkaloids, and flavonoids. The seed gum is composed of galactose and mannose [21]. The seeds also contain alkaloids such as trigonelline, gentianine, and calpain [26]. The bitter taste of fenugreek seeds is due to the presence of oils, steroidal saponins and alkaloids. It is rich in trace elements (Ca, Cu, Fe, Zn, K, Mg, P, Na, Mn), Vitamin A, B1 and C, sitosterol, n-alkanes, sesquiterpenes and cholesterol [27,28].

The scent dynamic mixes dependent on the fenugreek fragrance identification with the assistance of Gas Chromatograph and these incorporate the olfactometry diacetyl, 1-Octene-3-one, sotolon, acidic corrosive; 3-Isobutyl-2-methoxypyrazine, butanoic corrosive, isovaleric corrosive, 3-isopropyl-2-methoxypyrazine, caproic corrosive, eugenol, 3-Amino-4,5-dimethyl-3, linalool, (Z)-1,5-Octadiene-3-one, 4-dihydro-2 (5H)-Furanone with trademark smell of rich like roasty/natural, metallic, sharp, paprika like, sweat-soaked/foul, flowery, smelly, zesty respectively. The extract containing trigonelline and trigonellinic acid is useful as a hair growth stimulant (Table 2).

Table 1. Major Nutrients, carbohydrates, vitamins and mineral properties of fenugreek seed (value /10g)

Major Nutrient	Value g/10 g	Vitamins	Value mg/10 g
Total Calories	32.30	Vitamin A	6.00
Calories from Fat	5.40	Vitamin B ₁	0.04
Total Fat	0.60-0.79	Vitamin B ₂	0.03
Protein	2.30	Vitamin C	0.30
Total Carbohydrate	5.80	Thiamin	0.03
Water	0.24-0.88	Riboflavin	0.04
Dietary Fiber	2.40	Niacin	0.16
Mucilage	86.10	Vitamin B ₆	0.06
Fat, Lipids & Fatty Acids	Value g/10 g	Folate (mcg)	5.70
Saturated fat	150.00	β -carotene	0.96
Total fatty acid content (%)	4.50 - 7.00	Minerals	Value mg/10 g
Palmitic acid	53.00	Calcium (Ca)	17.60
Oleic acid	73.00 – 117.00	Iron (Fe)	3.30
Linoleic acid	225.00 – 360.00	Magnesium (Mg)	19.00
Linolenic acid	110.00 -175.00	Phosphorus (P)	30.00
Steric acid	27.00	Potassium (K)	77.00
Free fatty acids	0.01	Sodium (Na)	6.70
Monoacylglycerols	0.01	Zinc (Zn)	0.25

Source: USDA Nutrient Database. mg/10g- mili gram per 10 gram; g/10g-gram per 10 gram

In recent years there has been a marked increase in medicinal plants worldwide. The burgeoning problem due to modern medical treatments, methods, and harmful side effects of those drugs has attracted the general public attention, especially medical scholars. In a country where a large majority of people live in villages, live below the poverty line, and cannot afford expensive medical care, it poses a severe threat to the health of ordinary men [29].

6. PHARMACOLOGICAL ACTIVITIES FOR HUMAN HEALTHCARE

In the history of fenugreek, it has been used for various reasons by ancient Egyptians and played

a role in both herbal medicine and religious practices. In Greece, fenugreek is still used as animal feed. Thus, the Latin name for fenugreek *Trigonella foenum-graecum* means 'hay' in Greek. Thousands of years ago, herbaceous plants were highly valued in the cultures of Egypt, China, India, Arab countries, the Roman Empire, and Greece. Because of its role in health care, the functional food industry has classified it as a potential dietary supplement. Much research has been done on therapeutic effects and compound identification. The pharmacological actions of fenugreek, such as antioxidant, anti-cancer, gastroprotective, antidiabetes antirheumatic, etc., are discussed below (Tables 2 & 3).

Table 2. Proximate composition of fenugreek and their healthcare benefits

Chemical constituents	Effective against	Reference
Trimethylamine, Neurin, Trigonelline, Choline, Gentianine, Carpaine and Betain vitexin, Tricin, Quercetin, naringenin, rutin, isovitexin, Antioxidant	Hypoglycemic, antinociceptive, anti-inflammatory santi-inflammatory	[52,53] [52,53,54,55,56]
Isoleucine, 4-Hydroxyisoleucine, Histidine, Leucine, lysine, L-tryptophan, Arginine	Diabetes Mellitus,	[57]
diosgenin Graecunins, fenugrin B, fenugreekine, trigofenosides A–G Gum, neutral detergent fiber	Diabetes, hypercholesterolemia Digestion, constipation, heart problems, insulin response	[58,59,60] [61,62,63,64]
Triacylglycerols, tocopherols, sterols, free fatty acids.	Low Cholesterol	[63]
Caryophyllene	Essential oil, used in food products, bioactive properties	[64]
Trichloroacetic acid, pentadecyl ester Phytol	Used in cosmetic treatments Used in antioxidant, anti-inflammatory, antibacterial, anti-asthma, anti-cancer, and personal care products to produce synthetic forms of vitamin E and vitamin K1, also used in sexual and urological disorders, bioactive	[65] [66,67,68]
Quinoline	Antiasthmatic, Antitussive, and Antibacterial agents are bioactive and also used for urinary disorders.	[69,70]
Tetradecane	Drugs for disorders of the nervous system, antioxidants, antibiotics, and antiseptics also used as food additives, bioactive properties	[71,72]
Diosgenin	Suppressed tumour mass and volume of cancer cells, Induced cytotoxicity; Diminished tumour growth; Inhibited proliferation, migration, and invasion	[73,74,75]
Protodioscin	Induced reduction of cell growth and increased apoptosis	[76]

Table 3. Doses and concentrations of fenugreek products and compounds

Compounds/products	Concentration	References
Diosgenin	1-10 μ M	[77]
4-Hydroxyisoleucine	5-200 μ M	[78, 79]
Trigonelline	40 mg/kg (diet)	[80]
Fenugreek seed extract (FSE)	100 mg/kg	[81]
Fenugreek seed powder	9 g/kg (diet), 3%	[81,82]

6.1 Antioxidant Effects

Oxidative damage to proteins and lipids is caused by the overproduction of reactive oxygen species, leading to chronic degenerative diseases. Several studies have suggested fenugreek as a potential antioxidant [29]. The protective effects of fenugreek were demonstrated against lipid peroxidation and enzymatic antioxidants in cyclophosphamide-treated mice by assessing the extent of lipid peroxidation and the presence of antioxidants in the bladder of mice [30]. Fenugreek contains many essential nutrients, and this makes it a potent antioxidant. Supplementing the diet with fenugreek seeds decreased lipid peroxidation, increased glutathione and beta-carotene levels, and decreased alpha-tocopherol levels. This study shows that impaired free radical metabolism can be normalized by dietary fenugreek seed supplementation [31]. A study showed that fenugreek showed the greatest superoxide and radical scavenging effect and it was concluded that the antioxidant activity was related to high phenolic content in fenugreek [32].

6.2 Anti-diabetic Effect

Fenugreek seed powder solution remarkably improved lipid metabolism in patients with Type II diabetes. Fenugreek seeds may offer a new option for treating type II diabetes. Fenugreek seed contains about 0.1-0.5% 4-Hydroxyisoleucin that helps the stimulation of insulin secretion. Apart from that, the main saponin compound of fenugreek is diosgenin, which can inhibit cholesterol absorption and lower cholesterol levels in the liver. The extract showed significant activity against alloxan-induced diabetic conditions. With appropriate dosage, the 4-hydroxyisoleucine (amino acid) promotes insulin synthesis, hence controlling blood sugar level. The hypoglycemic effect of the extract was compared with that of the anti-diabetic glimepiride. The extract 4 mg/kg single dose showed significant activity against alloxan-induced diabetic conditions. The most effective dose recognized was 1 g/kg, lower than standard

anti-diabetic drugs [33]. In another study, a fenugreek seed was included in the diet of normal and diabetic hypercholesterolemic dogs. The defatted fraction contained dietary fibre (54%) and steroidal saponins (5%) reduced plasma cholesterol levels, glucagon and blood glucose, from pretreatment levels in dogs in both groups. The cholesterol-lowering effect was replicated in rats. Administration of high-fibre fractions of fenugreek lowered critical parameters, including total cholesterol, triglycerides, and low-density lipoproteins in diabetic rats [21].

6.3 Anti-cancer Effect

Cancer is becoming one of the major diseases worldwide, and it is the second leading cause of death worldwide after cardiovascular disease. Conventional cancer treatments only add a few months or years to a patient's life and cause significant side effects. Fruits and vegetables act as an alternative medicine and their active constituents are suggested for cancer treatment. Several plant-based anti-cancer drugs like vinca alkaloids (vinblastine and vincristine), epipodophyllotoxins (etoposide and teniposide), taxanes (paclitaxel and docetaxel), camptothecin derivatives (camptothecin and camptothecins) are available in the market. It is reported that the incidence of colon cancer can be reduced by including fenugreek seed powder in a diet [34]. A study revealed that fenugreek seed extract showed cytotoxic activity to inhibit the development of MCF-7 cells. Such effects have not been observed in liver cancer cell lines. These results highlight that the effects of fenugreek seed extract directly depend on the cancer cell type [35]. In another study, different normal and cancer cell lines were exposed to different concentrations of fenugreek extract (100 μ g/ml, 200 μ g/ml, and 300 μ g/ml) and different time points (0, 24, 48, 72, and 96 hours). Selective cytotoxic effects of fenugreek extract have been observed to act against various cancer cell lines, including T-cell lymphoma under *in vitro* conditions [36]. Some researchers proved that dietary intake of fenugreek seeds containing diosgenin could prevent

azoxy methane-induced colon cancer during the initiation and acceleration phases. These *in vitro* experiments on cancer cells demonstrated that diosgenin present in fenugreek seeds inhibited cell proliferation and induced apoptosis in the human colon cancer cell line HT-29 in a dose-dependent manner [37].

6.4 Anti-inflammatory and Anti-arthritic Effect

The body's defence mechanism, inflammation, is a complex biological response of vascular tissue to potentially harmful external or internal stimuli, including pathogens, chemicals, foreign substances and xenobiotics [38]. The lysosomal enzymes cause inflammatory disease release of the lysosomal enzyme, which stimulates prostaglandin synthesis. Petroleum ether was for extraction from fenugreek seed powder by cold maceration. This fenugreek seed petroleum ether extract (FSPEE) was analyzed by gas-liquid chromatography (GLC). Treatment with 0.5 ml/kg FSPEE reduced carrageenan- and formaldehyde-induced paw inflammation by 37% and 85%. A biphasic increase followed by a decrease in paw volume has been observed in CFA-induced arthritis. In rats treated with FSPEE (0, 5 ml/kg), cotton pellet weight was reduced by 42.5% and reduction in higher SGPT and ALP activities in serum and liver. Therefore, due to the presence of linoleic acid and linoleic acid, the fenugreek seed petroleum ether extract exhibits significant anti-inflammatory and anti-arthritic effects [39]. A study recorded an anti-inflammatory effect of fenugreek extract, which showed that rats were treated with a single dose (100 or 200 mg/kg). Inhibition of inflammatory swelling was 45% and 62% in the lower and higher dose groups, respectively, compared to 100% in untreated rats [39].

6.5 Antigastric Effect

Gastroprotective means any substances that have the ability for the prevent or ameliorate of injury to the gastrointestinal tract. The role of fenugreek seeds and omeprazole in ethanol-induced gastric ulcers has been studied. The results showed that the aqueous extracts and gel fractions isolated from the fenugreek seeds exhibited significant antiulcer protective effects. The cytoprotective effects of seeds appeared to be due not only to their antisecretory activity but also to their effects on mucosal glycoproteins [40].

6.6 Hepatoprotective and Nephroprotective Effects

Drugs or substances that have a beneficial effect on the liver are understood to be hepatoprotective. Sodium nitrite is one of the synthetic food additives that humans are constantly exposed to. Increased consumption of nitrite-treated products is directly related to the development of tissue damage, hepatotoxicity and nephrotoxicity. Hepatotoxicity and chronic liver injury from various causes are severe metabolic disorders that affect people of all ages [41]. A study was conducted on 40 rats in which rats were randomly assigned (10 per group) to control (physiological saline solution), fenugreek (150 mg/kg/day), NaNO₂ (80 mg/kg/day), and NaNO₂+fenugreek (80 mg/kg/day + 150 mg/kg/day) groups. This group was offered fenugreek seed extract two hours before NaNO₂. After three months, the rats were decapitated and blood, kidney and liver tissues were removed for the study. Results suggested that exogenous fenugreek mitigates NaNO₂-administration-induced hepatotoxicity and nephrotoxicity. Fenugreek extract exerted antioxidant and anti-inflammatory effects and played an essential role in preventing liver and kidney damage from chronic administration of NaNO₂ [42]. This study demonstrated fenugreek seeds' healing and protective effects against cisplatin-induced liver injury in rats at 5% and 10% levels of fenugreek supplementation. Another study investigated the defensive effect of aqueous extract of fenugreek against carbon tetrachloride-induced hepatotoxicity in rats. A 4% aqueous fenugreek extract enhanced the detrimental effects of carbon tetrachloride and normalized all biomarkers tested to control values. It has been concluded that it protects against liver damage caused by carbon tetrachloride. Results confirmed that carbon tetrachloride increased the antioxidant activity of serum transaminases, liver lipid peroxidation and liver enzymes [43].

6.7 Antibacterial and Antifungal Effects

Researchers have studied the various plants for antibacterial properties for decades in hopes of developing new treatments, including fenugreek. Botanical and microbial species use and the extraction method, influence the antimicrobial potency and severity [29]. Fenugreek seeds are known to be nutritious and good for digestion and it is commonly used as a spice to flavour curries and other recipes. In Ayurveda, the traditional Indian medicine, fenugreek seeds were used to

treat many gastrointestinal ailments. In a study, different concentrations of 125, 250, 500, and 1,000 $\mu\text{g/ml}$ fenugreek seed ethanol extract were used against bacteria. Most of the bacteria (*Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Proteus mirabilis*, *Escherichia coli* and *Vibrio parahaemolyticus*) were inhibited at a minimum inhibitory concentration of 50 $\mu\text{g/ml}$. These results are significant due to the small amount of fenugreek seed extract required to inhibit bacterial growth [35]. The fenugreek seed extract was investigated and found effective in controlling the growth of *Aspergillus flavus*, *Trichoderma viridiae* and *Trichophyton rubrum* when applied at 250 $\mu\text{g mL}^{-1}$ [30].

6.8 Infertility Effect

No studies have confirmed whether fenugreek seeds cause fertility or infertility. Several studies on fenugreek seeds have shown that fenugreek seeds have anti-implantation and anti-fertility effects and have abortion effects in rats. The dose of Fenugreek seeds of 1 g/kg/day orally reduces male fertility by lowering testosterone and sperm concentrations and inhibiting sperm mass and individual mortality [44]. The presence of steroidal oestrogen-like saponins may hurt serum testosterone and semen quality that ultimately acts as an anti-fertility agent [45]. High dietary levels of fenugreek significantly reduce fertility in both male and female rabbits, reduce testis weight, and cause obvious damage to seminiferous tubules and interstitial tissue [46]. The extract also showed weak estrogenic activity in immature ovariectomized rats. All observations indicate that fenugreek saponin extract has anti-fertility effects. Saponin extracts were found to be more effective in anti-implantation and abortion activities. The extract also showed weak estrogenic activity in immature ovariectomized rats. It shows that the substance has an anti-fertility effect [47]. The estrogenic effects exhibited by saponin extracts from fenugreek may be due to the presence of diosgenin, yamogenin-3 and tigogenin [48].

6.9 Weight Loss

Repeated administration of fenugreek seed extract slightly but significantly reduced dietary fat intake in healthy overweight individuals. These botanical components justified that fenugreek seed aqueous extracts effectively inhibited fat accumulation and ameliorated dyslipidemia in obese rats on a high-fat diet (HFD). This happened due to increased insulin sensitivity, glucose and lipid metabolism,

antioxidant defence and downregulation of lipogenic enzymes [49]. Significant weight loss was observed in mice with a high concentration of fenugreek (1%). Fenugreek seed extract effectively reduces body and adipose tissue weight dose-dependently [50]. In another study, fenugreek seed proteins, galactomannans, and polyphenols have been previously reported to modulate dyslipidemia in obese and diabetic rodents [51]. Further studies are needed to investigate fenugreek and other anti-obesity mechanisms.

7. FENUGREEK AS FODDER FOR ANIMALS

Animal components are the backbone of Indian agriculture in arid and semi-arid regions of the country. Round the year availability of diverse green fodder is paramount to providing balanced nutrition and fiber-rich diet to animals (small, medium and large) for their better growth, development and yields (milk, dung, wool). Feeding fenugreek grain to cattle has been a common practice for ages in India to avoid stomach-related problems and improve animal milk quality. Based on a three-week study with dairy cows, has shown that dietary fenugreek grains (20% of dry diet matter) increased milk yield and concentration of most of the functional fatty acids while reducing the blood cholesterol concentration in milk without altering milk flavour or taste [83]. One study concluded that fenugreek fed to crossbred Damascus goats resulted in significantly higher serum prolactin and increased milk production compared to the control. This may be mediated by prolactin hormone stimulation. Goats fed 60 g/day of fenugreek seed powder had significantly higher milk yields than the control group (1236 vs 1093 ml/day) of goat [84].

Fenugreek constitutes high-quality nutritious feed for dairy cattle, improving livestock's health status [15] because of a good combination of bioactive components (flavonoids, polyphenols, proanthocyanidins). Fenugreek grains and shoots could be an alternative option to enrich animal feed's biological and nutritional value in case of the unavailability of regular green fodder like lucerne and berseem. Fenugreek, a legume with a very good composition of biologically active components, represents differently in various anatomic parts. Supplementing animal feed with fenugreek could prevent birth abortion and better management of babies during lactation in cows, buffaloes and other animal models (Table 4).

Table 4. Significance of fenugreek as fodder for animals

Animal used for the study	Beneficial role	References
Buffaloes	A feed rich in ground fenugreek seed (GFS) enhanced total dry matter (13.17–14.00 kg/day) and milk output (7.34–8.01 kg/day) in Anatolian water buffaloes (AWBs). Adding fenugreek seeds (1.5%) to the feed of Surti buffaloes increased milk output while improving sensory attributes.	[85,86]
Dairy cow	Fenugreek seed feeding increased milk's functional fatty acid profile, decreased blood cholesterol levels (4% more than the control diet), and reduced cholesterol levels in milk (1.48 vs. 1.83 mg/g milk lipid).	[83]
Goat	A decrease in goat milk fat owing to fenugreek seed supplementation at very high doses. Interestingly, there was a positive trend in the flavour and palatability of boiling milk, indicating that fenugreek supplementation had a beneficial impact on milk quality. Fenugreek supplementation improves milk output, plasma constituents and growth hormone levels in goats	[87,84]
Rabbits	Fenugreek seed gum (FSG) may be considered a prebiotic. It is based on its resistance to pre-caecal digestion and favourable influence on the caecal bacteria population. Combining fenugreek seeds and probiotics enhanced growth performance, nutrient digestibility, and serum hormones without affecting carcass parameters in either male or female-developing rabbits.	[88,89]
Broiler chicken	The addition of seeds to the diet raised plasma concentrations of total protein and globulin. Supplementation of diets with a mixture of fenugreek and black cumin improved body weight gain, FCR, Hb, PCV, serum proteins, and serum cholesterol levels. 1 percent fenugreek powder for 38, 15 and 42 days produced significantly higher BW and BWG than the control groups of chicks. Fenugreek seed extract (FSE) is replace antibiotics to improve broiler health and growth and it increases chicken production while ensuring chicken quality and health. The optimal concentration of 100 mg/kg for FSE has been initially found.	[81,90,91, 92]

mg/g; Kg/day-kilogram per day; BW-Body weight; BWG-body weight gain; FCR-feed consumption ratio; Hb-Hemoglobin; PCV-packed cell volume

8. PRECAUTIONS

As per Ayurveda, fenugreek seeds may not be appropriate for individuals with the 'Pitta' body type since it is probably builds the 'Pitta' side effects, heat sensation and heat creation in the body. Because of this impact, a few people experience expanded perspiring after consuming fenugreek seeds. The most common side effects of fenugreek are loose stools when it is

consumed for the first time, excessive use can cause runs, perspiring problems, a headache, apprehension, quick heartbeat, insecurity, untimely labour and premature delivery, stomach agitation and sometimes swelling. This may cause allergy symptoms like skin rash, tingling, swelling of face, tongue, or throat swelling, dizziness and trouble breathing. Fenugreek seeds have hot potency and produce a heat sensation in the body. Hence, Fenugreek seeds

contradict bleeding issues since seeds may promote bleeding. Then again, fenugreek leaves have inverse activity because of their cool intensity. It leaves also used in Pitta Dosha and reduce bleeding, as indicated in Ayurveda. Hence other than normal cooking diets, it is very much necessary to consume this medicinal plant with the advice of some Ayurvedacharya or doctor for its better results without any side effects.

9. CONCLUSION

The properties of plants in the Indian tradition were discovered differently than in Western science. This is described in traditional textbooks as "Dravya-Guna-Shastra" (the science of the properties of materials). The challenge for Indian scientists is to build cross-cultural bridges between Dravya-guna-shastra and modern pharmacology. Such bridges can only be built if mutual respect exists and scientists feel they own indigenous knowledge and modern science. Though there are sufficient research data and literature available for fenugreek use and its benefits to prove this crop one of the best medicinal, easily indigenously available wonder herb and multipurpose crop with traditional values in pharmaceutical, nutraceutical, feed, fodder and industrial applications but still there exists a big gap to fill by the knowledge and for scientific research to gain the attention of the world to this medicinal seed spice crop. This low volume high-value, nutritionally rich, nutraceutical important, multipurpose plant has a huge potential and applications in the cuisines and feed, medicines, cosmetics and nutraceuticals and pharmaceutical industries. This review will be for end users to improve their health, improved immunity in human beings and higher milk production with good health in cattle. Looking to its diverse effects and variable constituents, it is very much required to do screening of available germplasm for its medicinal and aromatic contents available in variable amounts. It requires some in-depth scientific experiments to fix its medicinal products as readily available and safe consumable form so that the products may reach every end user and be used efficiently. This plant contains medicinal alkaloids, steroidal compounds and saponins, and many uses of this plant are mentioned in traditional medicine. Fenugreek primarily addresses non-communicable diseases, which are currently prevalent worldwide. It can be used as a preventive measure rather than solely for treatment. As a

common spice, fenugreek is readily available and can be easily incorporated into a busy lifestyle. The clinical use of fenugreek can be attributed to the abundant chemical constituents that fenugreek possesses. Since bioactive compounds have a promising future in disease-curing natural products, further research and research will be conducted to isolate bioactive compounds from crude extracts for drug development.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

The authors acknowledge the Director ICAR-National Research Centre on Seed Spices, Ajmer, Rajasthan for providing opportunity and motivation to work at NRCSS, Ajmer.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Blank I, Lin J, Devaud S, Fumeaux R, Fay LB. The principal flavour components of fenugreek (*Trigonella foenum graecum*). Risch SJ, Chi TH (Eds.), Spices: Flavour Chemistry and Antioxidant Properties, ACS, (Washington, DC), Publisher: American Chemical Society. 1997;12-28.
2. Flammang A, Cifone M, Erexson G, Stankowski L. Genotoxicity testing of a fenugreek extract. Food Chem. Toxicol. 2004;11:1769–1775.
3. De Candolle A, Origin of Cultivated Plants. (New York, Hafner). 1964;468.
4. Rao PG, Sriramulu M. Physiological characterisation of a spice (*Coriandrum sativum*) and a condiment (*Trigonella foenum-graecum*) during vegetative and reproductive stages. Curr. Sci. 1977; 46:615–617.
5. Duke JA, Reed CF, Weder P. *Trigonella foenum-graecum* L. In: Handbook of legumes of world economic importance. edited by J A Duke (Plenum Press, New York); 1981.

6. Al-Habori M, Raman A. Antidiabetic and hypocholesterolaemic effects of fenugreek. *Phytother Res.* 1998;12:233–242.
7. Pandey AK, Tripathi SC, Singh HN, Singh SB. Fungitoxicity of *Trigonella foenum-graecum* L. against some sugarcane pathogens. *Sugar. Cane.* 1993; 3:15–18.
8. Devi LS. Evaluation of green manures against the pigeon pea cyst nematode *Heterodera cajani*. *Natl. Acad. Sci. Lett.* 1997;20:1–2.
9. Nutman PS. Alternative sources of nitrogen for crops. *J. Royal Soc. Agric. England.* 1976;137:86–94.
10. Desperrier N, Baccou J C, Sauvaire Y. Nitrogen fixation and nitrate assimilation in field grown fenugreek. *Plant Soil.* 1986; 92:189–199.
11. Whistler R. Introduction to industrial gums. In: *Industrial gums*, edited by Whistler R L, BeMiller J N, Academic Press, (New York); 1993.
12. Garti N, Madar Z, Aserin A, Sternheim B. Fenugreek galactomannans as food emulsifiers. *Food Sci. Technol.* 1997; 30:305–311.
13. Godara AS, Singh R, Chouhan GS, Nepalia V. Yield and economics of fenugreek (*Trigonella foenum-graecum* L.) as influenced by fertility levels, biofertilizers and brassinosteroid. *Legume Res.* 2017; 40(1):165-169.
DOI :10.18805/lr.v0iOF.11192
14. Żuk-Gołaszewska K, Wierzbowska J. Fenugreek: productivity, nutritional value and uses. *J. Elem.* 2017;22(3):1067-1080.
DOI: 10.5601/jelem.2017.22.1.1396.
15. Solorio-Sánchez F, Solorio-Sánchez B, Basu SK, Casanova-Lugo F, Sarabia-Salgado L, Ku-Vera J, Cetzal-Ix W. Opportunities to grow annual forage legume fenugreek (*Trigonella foenum-graecum* L.) under mexican sylvopastoral system, *Am. J. Social Issues Humanities.* 2014;86–95.
16. Choudhary S, Singh R, Meena RS, Jethra G. Secondary and tertiary structure prediction of fenugreek (*Trigonella foenum-graecum*) protein. *Legume Res.* 2016; 39(1):48-51.
17. Choudhary S, Meena RS, Singh R, Vishal MK, Choudhary V, Panwar A. Assesment of genetic Diversity among Indian fenugreek (*Trigonella foenum-graecum* L.) varieties using morphological and RAPD markers. *Legume Res.* 2013;36(4):289-298.
18. Ali SF, Lal G, Aiswath OP, Chahar OP, Choudhary S, Mathews C, Anwer MM. Possibilities and potential of Rhizobial inoculants in organic production of fenugreek in arid and semiarid region of Rajasthan. *Intl. J. Seed Spices.* 2012; 2(1):39-45.
19. Bhandari DC. Agrobiodiversity conservation for sustainable agriculture development. Proceedings of IPPC-2018, International Congress on “Plant Based Natural Products: Phytocosmetics, Phytotherapeutics and Phytonutraceuticals” (Udaipur, Rajasthan, India). 2018;3-4.
20. Bieńkowski T, Żuk-Gołaszewska K, Kurowski T, Gołaszewski J. Agrotechnical indicators for *Trigonella foenum-gracum* L. production in the environmental conditions of northeastern Europe. *Turk. J. Field Crops.* 2016;21(1):16-28.
21. Snehlata HS, Payal DR. Fenugreek (*Trigonella foenum-graecum* L.): An overview. *Intl. J. Cur. Pharma Rev. and Res.* 2012;2:169–187.
22. Rao AV. *Herbal Cure for Common Diseases*, Fusion Books, (New Delhi); 2003.
23. Yadav S, Sehgal S. Effect of home processing and storage on ascorbic acid and β-carotene content of bathua (*Chenopodium album*) and fenugreek (*Trigonella foenum-graecum*) leaves. *Plant Food Hum. Nutr.* 1997;50:239-247.
24. Bukhari SB, Muhammad IB, Shahabuddin M. Antioxidant activity from the extract of fenugreek seeds. *Pak. J. Anal. Environ. Chem.* 2008;9(2):78-83.
25. Zandi P, Basu SK, Cetzal-Ix W, Kordrostami M, Chalaras SK, Khatibai L B. Fenugreek (*Trigonella foenum-graecum* L.): An important medicinal and aromatic crop, In: *Active Ingredients from aromatic and medicinal plants.* edited by H El-Shemy. 2017;207–224.
26. Mandgary A, Pournamdari M, Sharififar F, Pournourmohammadi S, Fardiar R, Shooli S. Alkaloid and flavonoid rich fractions of fenugreek seeds (*Trigonella foenum-graecum* L.) with antinociceptiv and anti-inflammatory effects. *Food Chemical Toxicol.* 2012;50(7):2503-2507.
27. Bhatia K, Kaur M, Atif F, AliM, Rehman H, Rahman S. Aqueous extract of ameliorates additive urotoxicity of buthionine sulfoximine and cyclophosphamide in mice. *Food Chem. Toxicol.* 2006;44:1744-1750.

28. Naidu MM, Shyamala BN, Naik PJ, Sulochanamma G, Srinivas P. Chemical composition and antioxidant activity of the husk and endosperm of fenugreek seeds. *Food Sci. Technol.* 2010;44:451-456.
29. Mandal S, DebMandal M. Fenugreek (*Trigonella foenum-graecum* L.) oils, In: In essential oils in food preservation, flavor and safety, edited by V. R. Preedy, (Academic Press), 2016;421–429.
30. Premanath R, Sudisha J, Devi NL, Aradhya SM. Antibacterial and antioxidant activities of fenugreek (*Trigonella foenum-graecum* L.) leaves. *Res. J. Med. Plant.* 2011;5:695–705.
31. Tejaswini BA, Sireesha C, Kaladhar K, Rao DG, Surekha C. *In vitro* antioxidant and antimicrobial activities of crude extracts of *Trigonella foenum-graecum* seeds. *Asian J. Chem.* 2012;24:5019–5022.
32. Shah MA, Mir PS. Effect of dietary fenugreek seed on dairy cow performance and milk characteristics. *Can. J. Anim. Sci.* 2004;84:725–729.
33. Shankar D. Garden cure, FRIHT, The Hindu, (13/12/2003) (Banglore); 2013.
34. Al-Janabi, Abdul K, Ahmed F. Feeding effects of fenugreek seeds (*Trigonella foenum-graecum*) on lactation performance, some serum constituents and prolactin hormone level in Damascus crossbred goats. *Diyala Agric. Sci. J.* 2012;4(1):1–8.
35. Alamer MA, Basiouni GF. Feeding effects of fenugreek seeds (*Trigonella foenum-graecum*) on lactation performance, some plasma constituents and growth hormone level in goats. *Pak. J. Biol. Sci.* 2005;8(11):1553-1556.
36. Pride of India, A Glimpse into India's Scientific Heritage, Samskrita Bharti, (New Delhi), ISBN 81-8727627-4. 2006;208.
37. Sowmya P, Rajyalakshmi P. Hypocholesterolemic effect of germinated fenugreek seeds in human subjects. *Plant Food Hum Nutr.* 1999;53:359-365.
38. Bhutya RK. Ayurvedic medicinal plants of India. Vol. I, (Scientific publishers (India), Jodhpur); 2011.
39. Roberts KT. The potential of fenugreek (*Trigonella foenum-graecum*) as a functional food and nutraceutical and its effects on glycemia and lipidemia. *J. Med. Food.* 2011;14(12):1485-1489.
40. Zafar MI, Gao F. 4-hydroxyisolumicine: A potential new treatment for type 2 diabetes mellitus. *Biodrugs.* 2016;30:255-262.
41. Fatima T, Maqbool K, Hussain SZ. Potential health benefits of fenugreek. *Journal of medicinal plant studies.* 2018; 6(2):166-169.
42. Zia T, Hasnain SN, Hasan SK. Evaluation of the oral hypoglycaemic effect of *Trigonella foenum-graecum* L. (methi) in normal mice. *J. Ethnopharmacol.* 2001;75: 191-195.
43. Srivastava D, Rajiv J, Naidu MMM, Puranaik J, Srinivas P. Effect of fenugreek seed husk on the rheology and quality characteristics of muffins. *Food Nutr. Sci.* 2012;3:1473-1479.
44. Mathern JR, Raatz SK, Thomas W, Slavin J. Effect of fenugreek fiber on satiety, blood glucose and insulin response and energy intake in obese subjects. *Phytotherapy Res.* 2009;23(11):1543-1548.
45. Platel K, Srinivasan K. Influence of dietary spices and their active principles on pancreatic digestive enzymes in albino rats. *Nahrung.* 2000;44:42–46.
46. Chatterjee S, Variyar SP, Sharma A. Bioactive lipid constituents of fenugreek. *Food Chem.* 2010;119:349-353.
47. Polonskaya Y, Shramko VVS, Morozov SV, Chernyak EI, Chernyavsky AM, Ragino YI. Balance of fatty acids and their correlations with parameters of lipid metabolism and markers of inflammation in men with coronary atherosclerosis. *Bull Exp. Biol. Med.* 2017;164:33–35.
48. Sarikurkcu C, Ozer MS, Calli N, Popović-Djordjević J. Essential oil composition and antioxidant activity of endemic *Marrubium parviflorum* subsp. *Oligodon*. *Ind. Crops Prod.* 2018;119:209–213.
49. Ram M, Rao K. The GC-MS study of one ayurvedic preparation Katakakhadiradi Kashayam. *Intl. J. Pharm. Sci. Rev. Res.* 2016;39:216–224.
50. Pirbalouti A G, Mirbagheri H, Hamed B, Rahimi R. Antibacterial activity of the essential oils of myrtle leaves against *Erysipelothrix rhusiopathiae*. *Asian Pac. J. Trop. Biomed.* 2014;4:S505–S509.
51. Okiei W, Ogunlesi EO, Osibote EAS. Analysis of essential oil constituents in hydro-distillates of *Calotropis procera* (Ait.). *Res. Br. Res. J. Phytochem.* 2009;3:44–53.

52. Lee W, Woo ER, Lee DG. Phytol has antibacterial property by inducing oxidative stress response in *Pseudomonas aeruginosa*. *Free Radic. Res.* 2016;50:1309–1318.
53. Chabukswar AR, Kuchekar BS, Jagdale SC, Lokhande PD, Chabukswar VV, Shisodia SU, Mahabal R H, Londhe AM, Ojha NS. Synthesis and evaluation of analgesic, anti-asthmatic activity of (E)-1-(8-hydroxyquinolin-7-yl)-3-phenylprop-2-en-1 ones Synthesis, evaluation, analgesic anti-asthmatic activity of (E)-1-(8-hydroxyquinolin-7-yl)- 3-phenylprop-2-en-1 one. *Arab J. Chem.* 2016;9:704–712.
54. Ramesh E, Manian RDRS, Raghunathan R, Sainath S, Raghunathan M. Synthesis and antibacterial property of quinolines with potent DNA gyrase activity. *Bioorganic Med. Chem.* 2009;17:660–666.
55. Silvaa RO, Francisca NSC, Beatriz M, Sousaa S, Damascenoa RB. Phytol, a diterpene alcohol, inhibits the inflammatory response by reducing cytokine production and oxidative stress Phytol inhibits the inflammatory response. *Fundam Clin. Pharmacol.* 2014;28:455–464.
56. Wang D, Wang C, Pi X, Guo L, Wang Y, Li M, Feng Y, Lin Z, Hou W, Li E. Urinary volatile organic compounds as potential biomarkers for renal cell carcinoma. *Biomed Reports.* 2016;5:68–72.
57. Das S, Dey KK, Dey G, Pal I. Antineoplastic and apoptotic potential of traditional medicines thymoquinone and diosgenin in squamous cell carcinoma. *Plos One.* 2012;7:e46641.
58. Srinivasan S, Koduru S, Kumar R, Venguswamy G. Diosgenin targets Akt-mediated prosurvival signaling in human breast cancer cells. *Int. J. Cancer.* 2009;15:961–967.
59. Chen PS, Shih YW, Huang HC, Cheng HW. Diosgenin, a steroidal saponin, inhibits migration and invasion of human prostate cancer PC-3 cells by reducing matrix metalloproteinases expression. *Plos One.* 2006;6:e20164.
60. Hibasami H, Moteki H, Ishikawa K, Katsuzaki H. Protodioscin isolated from fenugreek (*Trigonella foenum-graecum* L.) induces cell death and morphological change indicative of apoptosis in leukemic cell line H-60, but not in gastric cancer cell line KATO III. *Intl. J. Mol. Med.* 2003;11:23–26.
61. Degirmencioglu T, Unal H, Ozbilgin S, Kuraloglu H. Effect of ground fenugreek seeds (*Trigonella foenum-graecum*) on feed consumption and milk performance in Anatolian water buffaloes. *Arch. Anim. Breed.* 2016;59:345–349.
62. Choubey M, Patel VR, Raval AP, Singh RR. Effect of fenugreek seed supplementation on lactation performance of Surti buffaloes. *Ind J Animal Sci.* 2018;88(3):322-325.
63. Shah M A, Mir P S. Effect of dietary fenugreek seed on dairy cow performance and milk characteristics. *Canadian J. Animal Sci.* 2004;84(4):725-729.
64. Balgees A, Atta Elmnan NM, Jame SA, Rahmatalla EO, Amasiab A, Mahala G. Effect of fenugreek (*Trigonella foenum-graecum*) seeds supplementation on feed intake, some metabolic hormones profile, milk yield and composition of nubian goats. *Res. J. Animal Sci.* 2013;7:1–5.
65. Alamer MA, Basiouni GF. Feeding effects of fenugreek seeds (*Trigonella foenum-graecum* L.) on lactation performance, some plasma constituents and growth hormone level in goats. *Pak. J. Biol. Sci.* 2005;8:1553–56.
66. Zemzmi J, Mabrouki S, Abdouli H, Najar T. Preliminary characterization of fenugreek seed gum for use as prebiotic in rabbits' nutrition, In: *Proceedings of the 20th International Symposium on Housing and Diseases of Rabbits, Fur Providing Animals and Pet Animals (Celle, Germany).* 2017;17-18.
67. Abdel-Wareth AAA, Fatma SOE, Zienhom SHI, Abdallah AG, Jayant L. Combined effects of fenugreek seeds and probiotics on growth performance, nutrient digestibility, carcass criteria, and serum hormones in growing rabbits. *Livestock Sci.* 2021;251:104616.
68. Al-Homidan IH, Ebeid TA, Al-Muzaini A, Abou-Emera OK, Mostafa MM, Fathi MM. Impact of dietary fenugreek, mungbean, and garden cress on growth performance, carcass traits, blood measurements, and immune response in broiler chickens. *Livestock Sci.* 2020;242:104318.
69. Yattoo MA, Sharma RK, Nazam K, Ankur R, Pathak K. Effect of fenugreek and black cumin seeds as feed additives on blood biochemical profile and performance of broilers. *Ind. J. Animal Nutrition.* 2012;29(2):174-178.

70. Weerasingha AS, Atapattu NM. Effects of fenugreek (*Trigonella foenum-graecum* L.) seed powder on growth performance, visceral organ weight, serum cholesterol levels and the nitrogen retention of broiler chicken. *Tropical Agric. Res.* 2013;24(3): 289-95.
71. Al-Timimi LAN. Antibacterial and anticancer activities of fenugreek seed extract. *Asian Pac J. Cancer Prev.* 2019; 1:20(12):3771-3776. DOI: 10.31557/APJCP.2019.20.12.3771.
72. Sujapandian R, Anuradha VV, Viswanathan P. Gastroprotective effect of fenugreek seeds (*Trigonella foenum-graecum*) on experimental gastric ulcer in rats. *J. Ethnopharmacol.* 2002;81:393–397.
73. Huang H, Wang X, Yang L, He W, Meng T, Zheng K, Xia X, Zhou Y, He J, Liu C, Zou S, Xiao D. The effects of fenugreek extract on growth performance, serum biochemical indexes, immunity and NF-κB signaling pathway in broiler. *Front. Vet. Sci.* 2022;9:882754. DOI: 10.3389/fvets.2022.882754
74. Geberemeskel GA, Debebe YG., Nguse, NA. Antidiabetic effect of fenugreek seed powder solution (*Trigonella foenum-graecum* L.) on hyperlipidemia in diabetic patients. *J. Diabetes Res.* 2019;5:2019: 8507453.
75. Vijayakumar MV, Pandey V, Mishra GC, Bhat MK. Hypolipidemic effect of fenugreek seeds is mediated through inhibition of fat accumulation and upregulation of LDL receptor. *Obesity.* 2010;18:667–674,
76. Kumar P, Bhandari U, Jamadagni S. Fenugreek seed extract inhibit fat accumulation and ameliorates dyslipidemia in high fat diet-induced obese rats. *Biomed. Res. Int.* 2014;606021. DOI: 10.1155/2014/606021.
77. Geetha M, Reddy SK, Krupanidhi AM, Muralikrishna KS, Patil N, Prashanth P. Effect of fenugreek on total body and organ weights: A study on mice. *Pharmacologyonline.* 2011;3:747-752.
78. Dande P, Patil S, Sateesh B. Evaluation of saponins from *Trigonella foenum-graecum* seeds for its antifertility activity. *Asian J. Pharm. Clin. Res.* 2012;5(3):154-157.
79. Mudathir AE, Shaddad SAI, Elsharif B, Algasem AAEA. Antifertility Effects of *Trigonella foenum-graecum* (fenugreek) ethanolic extract in male rats, cocks. *J. Pharm Biomed Sci.* 2013;32(32):1299-1304.
80. Sharma RD. Effect of fenugreek seeds and leaves on blood glucose and serum insulin responses in human subjects. *Nutr. Res.* 1986;6:1353-64.
81. Amira Kassem, Abdulwali Al-Aghbari, Molham AL-Habori T. Mohammed AlMamary. Evaluation of the potential antifertility effect of fenugreek seeds in male and female rabbits. *Contraception.* 2006;73:301-306.
82. Ahirwar D, Ahirwar B. Evaluation of antifertility activity of *Trigonella foenum-graecum* seeds. *Der Pharmacia Sinica.* 2010;1(3):33-39.
83. Snehlata H, Payal D. Fenugreek (*Trigonella foenum-graecum* L): An overview. *Intl. J. Curr. Pharm. Rev. Res.* 2012;2:169–187.
84. Uslu GA, Uslu H, Adali Y. Hepatoprotective and nephroprotective effects of *Trigonella foenum-graecum* L. (fenugreek) seed extract against sodium nitrate toxicity in rats. *Biomed. Res. Ther.* 2019;6:3142–3150.
85. Joglekar M, Mandal M, Murthy S. Comparative analysis of antioxidant and antibacterial properties of *Aegle marmelos*, *Coriandrum sativum* and *Trigonella foenum-graecum*. *Acta Biol. Indica.* 2012; 1:105–108.
86. Visuvanathan T, Than LTL, Stanslas J, Chew, SY. Vellasamy, S. Revisiting *Trigonella foenum-graecum* L.: pharmacology and therapeutic potentialities. *Plants.* 2022;11:1450
87. Mowla A, Alauddin M, Rahman MA, Ahmed K. Antihyperglycemic effect of *Trigonella foenum-graecum* (Fenugreek) seed extract in alloxan-induced diabetic rats and its use in diabetes mellitus: a brief qualitative phytochemical and acute toxicity test on the extract. *Afr. J. Tradit. Complement. Altern. Med.* 2009;6(3):255–261.
88. Bhatia K, Kaur M, Atif F, Ali M, Rehman H, Rahman S, Raisuddin S. Aqueous extract of *Trigonella foenum-graecum* L. ameliorates additive urotoxicity of buthionine sulfoximine and cyclophosphamide in mice. *Food Chem. Toxicol.* 2006;44(10):1744-50. DOI: 10.1016/j.fct.2006.05.013.
89. Al-Sultan SI, El-Bahr SM. Effect of Aqueous Extract of Fenugreek (*Trigonella foenum-graecum* L.) On selected

- biochemical and oxidative stress biomarkers in rats intoxicated with carbon tetrachloride. Intl. J. Pharmacol. 2015;11: 43-49.
90. Alsemari A., Alkhodairy F, Aldakan A. Al-Mohanna M, Bahoush E, Shinwari Z, Alaiya, A. The selective cytotoxic anti-cancer properties and proteomic analysis of *Trigonella foenum-graecum*. BMC Complement Altern. Med. 2014;14: 114.
Available:<https://doi.org/10.1186/1472-6882-14-114>.
91. Raju J, Patlolla JM, Swamy MV, Rao CV. Diosgenin, a steroid saponin of *Trigonella foenum-graecum* (Fenugreek), inhibits azoxymethane-induced aberrant crypt foci formation in F344 rats and induces apoptosis in HT-29 human colon cancer cells. Cancer Epidemiol Biomarkers Prev. 2004;13(8):1392–1398.
92. Pundarikakshudu K, Shah DH, Panchal AH, Bhavsar GC. Anti-inflammatory activity of fenugreek (*Trigonella foenum-graecum* L.) seed petroleum ether extract. Indian J. Pharmacol. 2016;48:441-4.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/123839>