



# **Combined Effect of Nitrogen and Phosphorus Fertilizer on the Activity of the Nitrate Reductase Enzyme in Different Wheat Cultivars**

**Pramod Kumar Sharma<sup>a++\*</sup>, Rekha Gurjar<sup>b++</sup>,  
Manoj Kumar<sup>c#</sup>, Devendra Kumar<sup>a</sup>  
and Amarnath Dabaria<sup>d++</sup>**

<sup>a</sup> Division of Soil Science and Agricultural Chemistry, College of Agriculture, RLBCAU, Jhansi, Uttar Pradesh-284003, India.

<sup>b</sup> Division of Soil Science and Agricultural Chemistry, SKNAU, Jobner, Jaipur, Rajasthan, India.

<sup>c</sup> Department of Agronomy, SKNAU, Jobner, Jaipur, Rajasthan, India.

<sup>d</sup> Division of Plant Breeding and Genetics, SKNAU, Jobner, Jaipur, Rajasthan, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i11667

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96793>

**Original Research Article**

**Received: 02/12/2022**

**Accepted: 05/02/2023**

**Published: 06/02/2023**

## **ABSTRACT**

Some characteristics of nitrogen metabolism were observed in the flag leaf on the main branch of wheat (*Triticum aestivum* L. and *Triticum durum* L.) In the current study, nine varieties were chosen and planted as test crop using split plot design and replicate at three times, with nutrient dose as the main plot and varieties as the sub plot treatment cultivated at three different soil nitrogen and

<sup>++</sup> M.Sc. Research Scholar;

<sup>#</sup> Ph.D. Research Scholar;

\*Corresponding author: E-mail: [pk258513@gmail.com](mailto:pk258513@gmail.com);

phosphorus levels (0, 60, and 120 kg/hac) and (0, 30, and 60 kg/hac). These nitrogen and phosphorus levels were handled using four different treatments, where T1 served as the control, T2 as the nitrogen and phosphorus dose at its optimal level, and T3 as the half-nitrogen and full-phosphate fertilizer doses. Fertilizer dosages of T4 that were half nitrogen and half phosphorus were studied. Flag leaf blades were seen to be engaged in NO<sub>3</sub> assimilation. The flag leaf blade had the highest nitrate reductase activity, free amino acid content, and soluble protein content among all the leaf blades. The activity of nitrate reductase was markedly increased by the addition of nitrogen to the soil. The presence of substrate-dependent enzyme activity was demonstrated in flag leaf tissue. A coincidental association between enzyme activity and the buildup of reduced nitrogen in the plant shown the rise in nitrate reductase activity in response to more nitrogen and the increase vegetative reduced nitrogen. A substantial positive connection was discovered between nitrate reductase activity (expressed as molesN/hac.each season) and grain nitrogen (kg N/hac) at maturity because the transfer of vegetative nitrogen to the grain was homogeneous across treatments. Additionally, there was a strong and positive correlation between seasonal nitrate reductase and grain yields (kg/hac.). In cultivars resistant to lodging, maintaining nitrate reductase activity during the reproductive period might boost grain protein production and avoid the decrease of grain protein percentage that is usually seen when grain yields are high.

**Keywords:** Flag leaf; lodging; wheat; phosphorus; nitrate reductase activity (NRA).

## 1. INTRODUCTION

Nitrogen is the most critical nutrient that plants get from the soil in terms of quantity. Lonhienne et al.,[1] Plant roots have long been known to take up nitrogen molecules with low molecular mass, such as ammonium, nitrate, and amino acids. In natural ecosystems, however, nitrogen is mostly found as proteins in the soil. This complex organic form of nitrogen thought to be unavailable to plants directly. Although roughly 80% of the nitrogen in the atmosphere is di nitrogen (nitrogen gas, N<sub>2</sub>), most living organisms cannot use this form of the element and converted into the useful form, such as ammonia. Bano and Iqbal [2], opined that Leguminous legumes' ability to fix di nitrogen into useful reactive nitrogen molecules has traditionally been utilized by humans and boosting soil fertility.

Proteins, Rubisco, nucleic acids, and chlorophyll all contain nitrogen as a structural component. N fertilisation plays an important agronomic management strategy for increasing crop productivity Astaneh et al. [3]. The supply of N in plants has a significant impact on the functional activity of the photosynthetic apparatus in leaves. Furthermore, it has been observed that effective N feeding has the capacity to mitigate drought stress effects by maintaining metabolic processes even at low tissue water potential Abidet al., [4]. Excess nitrogen allows many plants to grow aggressively with dark green coloured lush growth, but it also causes developmental alterations and may alter the

biology of plants such as a longer vegetative phase, delayed maturity, a longer plant life cycle, and enhanced succulence Dietz, et al.,[5].

The majority of nitrogen in the biosphere is in the form of N<sub>2</sub> in the atmosphere, which is useless to most of the plants species until it is "fixed" biologically or abiotically (by lightning or aurorae, or industrially). It is usually either absorbed and converted into biological N or nitrified into NO<sub>3</sub> once it is fixed into NH<sub>3</sub>. Ammonification is a process that converts organic nitrogen back into NH<sub>3</sub>. Nitrate can be transformed to N<sub>2</sub>O and N<sub>2</sub> through nitrification and denitrification, respectively. As a result of this N<sub>2</sub>O and N<sub>2</sub> production, ecosystems lose nitrogen while the atmospheric nitrogen stored in gains nitrogen.

The amount of nitrogen in a plant's tissues determines its growth. According to the N productivity idea, plants that develop in N-rich environments have higher internal N concentrations and a higher relative growth rate. Limited N supply causes low shoot growth, a high root–shoot ratio and decreased leaf growth in plants. As the leaves age more nitrogen is assigned to the highest leaves, where there is a greater demand for photosynthetic enzymes and chlorophyll and it is mobilized in the seeds Pilbeam [6].

Phosphorus (P) is an important nutrient for plant growth and productivity. It's content in plants ranges from 0.05 to 0.5 percent of the total dry weight of the plant. Despite the fact that the concentration of P in soil is 2000 times higher than in plants, its fixation in the form of

aluminium/iron or calcium/magnesium phosphates prevents plants from absorbing it Malhotra et al. [7]. Phosphorus is involved in a variety of biological functions, including membrane structure maintenance, biomolecule synthesis and the production of high-energy molecules. Nitrogen is the fourth most prevalent element in living and it is utilized to essential biological components including amino acids and nucleic acids Luo et al., [8]. All plants require the element phosphorus (P). Low amounts of phosphorus in the soil will cause plants to grow slowly, but for agricultural production reasons, the soil should have an adequate level of phosphorus to promote healthy plant growth Watson and Mullen [9].

Phosphorus (P) is a structural component of major bio molecules such as nucleic acids, sugar phosphates, adenosine triphosphates, and phospholipids, and an substitutable essential element for plant growth. Phosphorus (P) is one of the most important nutrients for plant growth and survival. It is essential for cellular bioenergetics and metabolic pathways within the plant body. The primary function of mineral fertilizers is to increase crop yield but the biggest impediment to realize known crop potential is the low or none use of fertilizers, notably P and N Irfan et al., [10].

In agricultural systems, the application of P to the soil is required to ensure plant productivity Simpson et al., [11]. Seed P reserves are rapidly mobilized and translocated to emerging root and shoot tissues after germination, as it is the sole P source available to sustain seedling growth. This P supply is then replenished by P uptake by the root system as it develops Julia et al., [12]. When root P acquisition is insufficient to meet the P requirement for new growth, biochemical, physiological, and morphological responses occur to improve tissue P economy and increase soil P acquisition White et al., [1]. Local and systemic signals involving gibberellins, auxin, cytokinins, ethylene, and strigolactones, as well as the translocation of regulatory RNAs and excess sucrose from the shoot to the root by the phloem, co-ordinate many of the responses of root tissues to P deprivation. Increased sucrose availability to the root in particular has been linked to increased P-uptake capability in P-deficient plants.

Phosphorus is the second most commonly limiting macronutrient for plant growth, behind N. It is a key plant macronutrient that accounts for around 0.2 percent of the dry weight of a plant.

Phosphorus is a component of important compounds including nucleic acids, phospholipids and ATP. Plants cannot thrive without a consistent supply of this nutrient. It is also involved in the regulation of metabolic pathways and the control of important enzyme processes Zhang et al. [13].

Kaure et al. [14] reported that the activities of nitrate reductase, nitrite reductase, glutamine synthetase, glutamate synthase, and glutamate dehydrogenase all increased as the nitrogen rate were increased, resulting in a rise in protein and amino acid content in all wheat genotypes. The amount of nitrogen and chlorophyll in the leaves decreased as the leaves grew. It was also shown that nitrogen assimilatory enzymes (nitrate reductase and glutamine synthetase) have a positive association with Nitrogen use efficiency (NUE) and nitrogen content, suggesting that these enzymes may be the rate limiting enzymes in nitrogen metabolism. Crops grown with standard nitrogen fertiliser had higher nitrogen and biomass contents, and the accompanying soils had more nitrification and denitrification genes in abundance Robinson et al. [15].

## 2. MATERIALS AND METHODS

The field study was carried out at the ICAR-IISS research farm in Bhopal, Madhya Pradesh, during the rabi season of 2020–21. It is classified as semi-arid and subtropical and has scorching summers and frigid winters. The Vindhyan Plateau Agroclimatic Zone includes Bhopal. About 1100 mm of rainfall falls on average each year, with the majority falling between July and September during the monsoon season. The average maximum summer temperature is 35–40°C, while the average winter minimum is 2–9°C. There are 36 plots in a block (9 variety x 4 fertiliser N and P treatments). Each plot is 2 m x 2 m in size.

The reduction of nitrate to nitrite was assayed *in vitro* by incubating the enzyme extract with KNO<sub>3</sub> in the presence of an electron donor (NADH). Nitrite was diazotized with sulphanilamide and then reacted with N-(1-naphthyl) ethylene diaminedihydrochloride (NEDD) to produce an azo dye which was measured spectrophotometrically at 540nm Nicholas and Nason [16]. The 0.2 gm leaf sample was taken and put into 0.2 M phosphate buffer solution and 0.2M KNO<sub>3</sub> (3 ML). The samples were kept in dark chamber for 1 hour for incubation of reaction. The reaction was terminated in boiling

water at 100 °C. Aliquot (0.5 ml) was taken to which 1 ml of 1N sulphanimide and 1ml of 0.02% NEDD was added and the volume was made up to 6 ml. The absorbance of the resultant solution (Pink colour) was measured against the blank at 540 nm wavelength. For the formation of standard curve potassium nitrite solution (0.01M) was formed with series of test tubes. The standard curve was prepared with diluted KNO<sub>2</sub> solution with series of test tubes and make up the volume in each to 2 ml with water. The enzyme activity was expressed as  $\mu\text{molKNO}_2/\text{h/g}$  fresh weight. The sampling for NRA was done at 50, 65 and 85 DAS (Days after sowing) on the morning hours.

## 2.1 Nine Varieties of Wheat

V<sub>1</sub> to V<sub>9</sub>=HI8663, HI8737, HI8713, HI1563, HI1544, HI1531, GW366, LOK1, NARMADA14 Recommended dose of fertilizer (RDF) @120:60:40 kg/ha of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O were supplied through Urea, SSP and MOP, respectively. 50% of N, 100% of P<sub>2</sub>O<sub>5</sub> and 100% of K<sub>2</sub>O of the respective treatments were applied as basal and rest 50% of the N dose was applied in two top dressings at 22 DAS and 45 DAS. The remaining N was top dressed in equal splits at 25 and 45 DAS, whereas 50% of the applied N was delivered as basal.

## 3. RESULTS AND DISCUSSION

### 3.1 Nitrate Reductase Activity (NRA)

At 50 DAS, there were non-significant differences in NRA observed between varieties of wheat and fertilizer treatments. The range of NRA was found in between 0.22(Narmada14) to 2.43  $\mu\text{moleKNO}_2/\text{h/g}$  (GW366) among all the treatments (Table 2 & Fig. 1). The mean NRA was higher in Normal dose treatment followed by reduced phosphatic fertilizer dose treatment, reduced nitrogen dose fertilizer treatment and lower in control plots. Among all the treatments, the highest magnitude of NRA was observed in GW366 (2.43  $\mu\text{mole KNO}_2/\text{h/g}$ ) followed by HI8663 (2.17  $\mu\text{moleKNO}_2/\text{h/g}$ ) in normal Dose fertilizer treatment and lower NRA was observed in NARMADA14 (0.22  $\mu\text{moleKNO}_2/\text{h/g}$ ) (Table 2 & Fig. 1).

At 65 DAS, the range of NRA was found in between 0.41(LOK1) to 1.73  $\mu\text{moleKNO}_2/\text{h/g}$  (HI1544) among all the treatments. The mean NRA was higher in Normal dose treatment

followed by reduced phosphatic fertilizer dose treatment and reduced nitrogen dose fertilizer treatment and lower in control plots (Table 2 & Fig. 2). Among the varieties grown in full dose of N & P treatment, the highest NRA was found in HI1544 (1.41  $\mu\text{moleKNO}_2/\text{h/g}$ ) and the lowest NRA was found in HI8713 (0.43  $\mu\text{moleKNO}_2/\text{h/g}$ ). Among the varieties grown in half dose of N fertilizer treatment the highest NRA was observed in HI1544 (1.54  $\mu\text{moleKNO}_2/\text{h/g}$ ) followed by HI1531 (1.44  $\mu\text{moleKNO}_2/\text{h/g}$ ) and among the varieties grown in half dose of P, the highest NRA was found in HI1544 (1.73  $\mu\text{moleKNO}_2/\text{h/g}$ ) followed by HI8713 (1.28  $\mu\text{moleKNO}_2/\text{h/g}$ ). Among all the treatments, the highest NRA was observed in HI1544 and lowest NRA was observed in LOK1 (0.41  $\mu\text{moleKNO}_2/\text{h/g}$ ) in reduced phosphorus dose fertilizer (Table 2 & Fig. 2).

At 85 DAS, there were significant differences in NRA observed between fertilizer treatment and varieties. The range of NRA was found in between 0.15(GW366) to 1.78  $\mu\text{moleKNO}_2/\text{h/g}$  (HI8713) among all the treatments (Table 2 & Fig. 3). The mean NRA was higher in Normal dose treatment followed by reduced phosphatic fertilizer dose treatment, reduced nitrogen dose fertilizer treatment and lower in control plots. Among the varieties grown in full dose of N & P treatment, the highest NRA was found in Narmada14 (1.57  $\mu\text{moleKNO}_2/\text{h/g}$ ) and the lowest NRA was found in HI1531 (0.33  $\mu\text{mole KNO}_2/\text{h/g}$ ). Among the varieties grown in half dose of N fertilizer treatment the highest NRA was observed in HI1563 (1.28  $\mu\text{moleKNO}_2/\text{h/g}$ ) followed by HI1544 (1.16  $\mu\text{moleKNO}_2/\text{h/g}$ ) and among the varieties grown in half dose of P, the highest NRA was found in HI8713 (1.78  $\mu\text{moleKNO}_2/\text{h/g}$ ) followed by HI1544 (1.42  $\mu\text{moleKNO}_2/\text{h/g}$ ). Among all the treatments, the highest NRA was observed in HI8713 (1.78  $\mu\text{moleKNO}_2/\text{h/g}$ ) grown in reduced P dose and lower NRA was observed in GW366 (0.15  $\mu\text{mole KNO}_2/\text{h/g}$ ) grown in control plots. Across all nutrient treatment leaf area in selected wheat varieties followed the following trends:-

HI8713 > LOK1 > HI8663 > HI1544 > HI8737 > NARMADA 14 > HI1563 > HI1531 > GW366 (Table 2 & Fig. 3).

The enzyme NR nitrate reduces nitrogen for protein metabolism in plant system. Nitrate is the principal source of nitrogen for wheat plant, wherein NRA is the rate limiting, and hence protein synthesis is mostly dependent on NR


activity. At all growth phases, NR activity was found to have a strong and positive relationship with grain protein. As a result, a favourable relationship between these two traits were inevitable Adavi et al. [17].

In this study, increased nitrate reductase enzyme activity was found in the HI1531 and HI8713 genotypes, and these genotypes also showed better grain production, indicating that the NRA and grain yield had a substantial relationship in the HI1531 and HI8713 genotypes. These

findings are in congruence with Fortunato et al. [18], wherein they found a highly substantial positive connection between NR activity and grain yield and grain protein. In wheat and triticale, Zhan et al. [19] revealed a strong positive association between NRA and grain yield. In addition Zhang et al. [20] discovered a substantial relationship between NRA in the top leaf during the tillering stage and grain yield and grain protein. As a result, NRA can be utilised to select genotypes with high grain protein levels.

**Table 1. Details of experiment**

1.	Design of experiment	Split plot design
2.	No. of replication	3
3.	No. of treatments (plots)	36
	Main plot	4
	Sub plot	9
4.	Net plot size	2m x 2m
5.	Gross plot size	2.5m x 2.5 m
6.	Row spacing	22.5 cm
7.	No. of rows per plot	9
8.	Crop	Wheat
9.	Plant population per plot	190
10.	Seed rate	100 kg ha <sup>-1</sup>
11.	Fertilizer dose (RDF)	120-60-40 kg ha <sup>-1</sup> of N, P and K
12.	Date of sowing	25 November 2020



**Fig. 1. Effect of N & P on NRA activity in leaf of wheat genotypes at 50 DAS**




Fig. 2. Effect of N & P on NRA activity in leaf of wheat genotypes at 65 DAS




Fig. 3. Effect of N & P on NRA activity in leaf of wheat genotypes at 85 DAS

Table 2. Effect of N & P on nitrate reductase enzyme activity ( $\mu\text{molKNO}_2/\text{h/g}$ ) of wheat genotype at 50, 65 and 85 DAS

	50 DAS					65 DAS					85 DAS				
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	Mean A	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	Mean A	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	Mean A
HI8663	0.90	2.17	0.82	1.34	1.31	0.64	0.86	0.93	0.64	0.77	1.03	0.87	0.41	1.51	0.95
HI8737	1.51	0.67	0.44	1.13	0.93	0.49	1.05	0.67	0.59	0.70	1.01	1.01	0.75	0.94	0.93
HI8713	0.50	0.66	0.88	1.26	0.82	0.48	0.43	0.29	1.28	0.62	1.32	1.38	0.83	1.78	1.33
HI1563,	0.63	1.35	0.50	0.68	0.79	1.08	1.28	1.06	1.17	1.15	0.34	1.56	1.28	0.16	0.84
HI1544	0.93	0.81	0.84	0.76	0.84	1.57	1.41	1.54	1.73	1.56	0.18	1.00	1.16	1.42	0.94
HI1531	1.46	0.79	1.79	1.01	1.26	1.10	1.19	1.44	0.81	1.13	1.15	0.33	1.11	0.75	0.83
GW366	0.65	2.43	0.57	0.87	1.13	0.59	0.84	0.70	0.47	0.65	0.15	1.40	0.84	0.91	0.82
LOK1	0.55	1.26	1.22	0.84	0.97	0.41	0.59	0.74	0.70	0.61	0.85	1.42	1.13	1.34	1.18
NARMADA14	0.76	0.22	0.90	0.25	0.53	0.80	0.75	0.79	0.81	0.79	0.31	1.57	0.79	0.87	0.89
Mean B	0.87	1.15	0.89	0.90		0.80	0.93	0.91	0.91		0.70	1.17	0.92	1.07	
<b>Factors</b>	<b>C.D.</b>	<b>SE(d)</b>			<b>SE(m)</b>	<b>C.D.</b>	<b>SE(d)</b>			<b>SE(m)</b>	<b>C.D.</b>	<b>SE(d)</b>			<b>SE(m)</b>
Factor(A)	NS	0.15			0.10	NS	0.08			0.06	0.06	0.02			0.01
Factor(B)	NS	0.41			0.29	0.36	0.18			0.13	0.13	0.07			0.05
Factor(B)at same level of A	NS	0.82			0.31	NS	0.35			0.17	0.27	0.13			0.04
Factor(A)at same level of B	NS	0.79			0.56	NS	0.34			0.24	0.26	0.12			0.09

T<sub>1</sub>=Control T<sub>2</sub>=100% (N+P+K) T<sub>3</sub>= 50% N+ 100% (P+K) T<sub>4</sub>=50%P+100%N+K

#### 4. SUMMARY AND CONCLUSIONS

At 50 DAS, GW366 had highest NRA content (2.43  $\mu\text{moleKNO}_2/\text{h/g}$ ) followed by HI8663 (2.17  $\mu\text{moleKNO}_2/\text{h/g}$ ) and at 65 and 85 DAS HI 1544 variety had highest average NRA content.

All of the investigated wheat varieties showed different levels of NRA activity, with some showing the highest concentrations at the early phases of flowering, others at the post-flowering stage, and still others at the vegetative stage.

The main findings are that (a) The rate of enzyme activity was significantly affected by the amount of nitrate present in the tissue. (b) Although not statistically, nitrate reductase activity was correlated with leaf protein content. (c) On a field scale, split dose of nitrogen proved successful in producing nitrate reductase. (d) Increases in protein (%) were associated with higher enzyme activity, which is achieved by dividing the nitrogen dose.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. White PJ, Veneklaas EJ. Nature and nurture: The importance of seed phosphorus content. *Plant and Soil*. 2012; 357:1-8.
2. Bano SA, Iqbal SM. Biological nitrogen fixation to improve plant growth and productivity. *Int. J. Agric. Innov. Res*. 2016;4:597-9.
3. Astaneh N, Bazrafshan F, Zare M, Amiri B, Bahrani A. Nano-fertilizer prevents environmental pollution and improves physiological traits of wheat grown under drought stress conditions. *Scientia Agropecuaria*. 2021;12(1):41-7.
4. Abid M, Tian Z, Ata-UI-Karim ST, Cui Y, Liu Y, Zahoor R, Jiang D, Dai T. Nitrogen nutrition improves the potential of wheat (*Triticumaestivum* L.) to alleviate the effects of drought stress during vegetative growth periods. *Frontiers in Plant Science*. 2016;7:981.
5. Dietz KJ, Zörb C, Geilfus CM. Drought and crop yield. *Plant Biology*. 2021;23(6):881-93.
6. Pilbeam DJ. The utilization of nitrogen by plants: a whole plant perspective. *Ann. Plant Rev. Online*. 2018;305-51.
7. Malhotra H, Sharma S, Pandey R. Phosphorus nutrition: Plant growth in response to deficiency and excess. *Plant nutrients and Abiotic Stress Tolerance*. 2018;171-90.
8. Luo C, Rodriguez-R LM, Johnston ER, Wu L, Cheng L, Xue K, Tu Q, Deng Y, He Z, Shi JZ, Yuan MM. Soil microbial community responses to a decade of warming as revealed by comparative metagenomics. *Applied and environmental microbiology*. 2014;80(5): 1777-86.
9. Watson M, Mullen R. Understanding soil tests for plant-available phosphorus. Unpublished. 2007;1755-315.
10. Khalofah A, Khan MI, Arif M, Hussain A, Ullah R, Irfan M, Mahpara S, Shah RU, Ansari MJ, Kintl A, Brtnicky M. Deep placement of nitrogen fertilizer improves yield, nitrogen use efficiency and economic returns of transplanted fine rice. *Plos One*. 2021;16(2):e0247529
11. Simpson RJ, Oberson A, Culvenor RA, Ryan MH, Veneklaas EJ, Lambers H, Lynch JP, Ryan PR, Delhaize E, Smith FA, Smith SE. Strategies and agronomic interventions to improve the phosphorus-use efficiency of farming systems. *Plant and Soil*. 2011;349:89-120.
12. Julia CC, Rose TJ, Pariasca-Tanaka J, Jeong K, Matsuda T, Wissuwa M. Phosphorus uptake commences at the earliest stages of seedling development in rice. *Journal of Experimental Botany*. 2018; 69(21):5233-40.
13. Zhang H, Hou J, Liu J, Zhang J, Song B, Xie C. The roles of starch metabolic pathways in the cold-induced sweetening process in potatoes. *Starch-Stärke*. 2017; 69(1-2):1600194.
14. Kaur B, Asthir B, Bains NS. Enzymatic efficiency and genotypic differences for nitrogen assimilation in wheat. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*. 2017;87:985-95.
15. Yeoh YK, Paungfoo-Lonhienne C, Dennis PG, Robinson N, Ragan MA, Schmidt S, Hugenholtz P. The core root microbiome of sugarcane cultivated under varying nitrogen fertilizer application. *Environmental microbiology*. 2016;18(5): 1338-51.
16. Nicholas DD, Nason A. [144] Determination of nitrate and nitrite.



17. Sathee L, Meena HS, Adavi SB, Jha SK. Nitric oxide-induced regulation of plant developmental processes and abiotic stress responses. *Plant Abiotic Stress Tolerance: Agronomic, Molecular and Biotechnological Approaches*. 2019;381-408.
18. Fortunato S, Nigro D, Paradiso A, Cucci G, Lacolla G, Trani R, Agrimi G, Blanco A, de Pinto MC, Gadaleta A. Nitrogen metabolism at tillering stage differently affects the grain yield and grain protein content in two Durum wheat cultivars. *Diversity*. 2019;11(10):186.
19. Chen Y, Zhang P, Wang L, Ma G, Li Z, Wang C. Interaction of nitrogen and phosphorus on wheat yield, N use efficiency and soil nitrate nitrogen distribution in the North China Plain. *International Journal of Plant Production*. 2020;14:415-26.
20. Zhang G, Liu S, Dong Y, Liao Y, Han J. A nitrogen fertilizer strategy for simultaneously increasing wheat grain yield and protein content: Mixed application of controlled-release urea and normal urea. *Field Crops Research*. 2022;277:108405.

---

© 2023 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/96793>